MONARCH NECTAR PLANTS





Left to right: Monarch on swamp milkweed, littleflower penstemon, and Rocky Mountain blazing star.

The Rocky Mountains region spans large sections of Idaho, Montana, Wyoming, Utah, Colorado, and New Mexico. It is characterized by some of the tallest mountains in the continental U.S. and is home to diverse forests, serpentine river canyons, and high alpine zones. A huge variety of plants and wildflowers, from tiny alpine blooms to large stands of rabbitbrush and goldenrod, can be found in this region. These plant communities support numerous pollinators and other wildlife, including summer-visiting monarch butterflies.

Each spring, monarchs leave overwintering sites in central Mexico and along the California coast and fan out across the North American landscape to breed and lay eggs on milkweed, the monarch's host plant. Several generations are likely produced during this time. In the fall, adults migrate back to the overwintering sites, where they generally remain in reproductive diapause until the spring, when the cycle begins again. Monarchs at overwintering sites in Mexico and California have declined dramatically since monitoring began in the late 1990s. Across their range in North America, monarchs are threatened by a variety of factors. Loss of milkweed from extensive herbicide use has been a major contributing factor, and habitat loss and degradation from other causes, natural disease and predation, climate change, and widespread insecticide use are probably also contributing to their decline. Because of the monarch's migratory life cycle,

it is important to protect and restore habitat across their entire range. Adult monarchs depend on diverse nectar sources for food during all stages of the year, from spring and summer breeding to fall migration and overwintering. Inadequate milkweed and nectar plant food sources at any point may impact the number of monarchs that successfully arrive at overwintering sites in the fall.

Providing milkweeds and other nectar-rich flowers that bloom where and when monarchs need them is one of the most significant actions you can take to support monarch butterfly populations. This guide features Rocky Mountain native plants that have documented monarch visitation, bloom during the times of year when monarchs are present in this region, are commercially available, and are known to be hardy. These species are well-suited for wildflower gardens, urban greenspaces, and farm field borders. Beyond supporting monarchs, many of these plants attract other nectar- and/or pollen-seeking butterflies, bees, moths, and hummingbirds. For a list of native plants that host butterflies and moths specific to your zip code see www.nwf.org/nativeplantfinder.

The species in this guide will be adaptable to growing conditions across most of the Rocky Mountains. Please consult regional floras or the Biota of North America's North American Plant Atlas (http://bonap.net/napa) for details on species' distributions in your area.










Bloom	Common Name	Scientific Name	Flower Color	Max. Height
	Forbs			(Feet)
Spring to Fall 1	Butterfly milkweed	Asclepias tuberosa	Orange/yellow	2
2	Showy milkweed	Asclepias speciosa	Pink/green/purple	4
3 Summer 4	Nettleleaf giant hyssop	Agastache urticifolia	Purple/red	5
	Tall fringed bluebells	Mertensia ciliata	Blue	3
5	Blanketflower	Gaillardia aristata	Red/yellow	3
6	Common sunflower	Helianthus annuus	Yellow	8
7	Dotted blazing star	Liatris punctata	Pink/purple	2
8	Flodman's thistle	Cirsium flodmanii	White/purple	3
9	Jerusalem artichoke	Helianthus tuberosus	Red/pink/yellow	10
10	Maximiliani sunflower	Helianthus maximiliani	Yellow/brown	10
11	Missouri goldenrod	Solidago missouriensis	Yellow	3
Summer to Fall 12	Rocky Mountain beeplant	Cleome serrulata	White/pink	4
13	Rocky Mountain blazing star	Liatris ligulistylis	Pink/purple	6
14 15	Spotted joe pye weed	Eutrochium maculatum	Pink/purple	6
	Sulphur-flower buckwheat	Eriogonum umbellatum	White/yellow	2
16	Swamp milkweed	Asclepias incarnata	Pink	4
17	Swamp verbena	Verbena hastata	Blue/purple	5
18	Western coneflower	Rudbeckia occidentalis	Yellow/green	6
19	Western goldentop	Euthamia occidentalis	Yellow	6
Fall 20	Canada goldenrod	Solidago canadensis	Yellow	5
	Shrubs			
Spring to Summer 21	Woods' rose	Rosa woodsii var. ultramontana	Pink	5
Summer 22	Littleflower penstemon	Penstemon procerus	Blue/purple	2
23 Summer to Fall 24	Rubber rabbitbrush	Ericameria nauseosa	Yellow	8
	Yellow rabbitbrush	Chrysothamnus viscidiflorus	Orange/yellow	3



Planting for Success

Monarch nectar and host plants often do best in open, sunny sites. You can attract more monarchs to your area by planting flowers in single species clumps and choosing a variety of plants that have overlapping and sequential bloom periods. Monarchs can be present from July through October in the Rocky Mountains, although this can vary depending on your elevation.

Why Plant Native?

Although monarchs use a variety of nectar plant species, including exotic invasives such as dame's rocket and yellow toadflax, we recommend planting native species. Native plants are often more beneficial to ecosystems, are adapted to local soils and climates, and help promote biological diversity. They can also be easier to maintain in the landscape, once established.

Tropical milkweed is a non-native plant that is widely available in nurseries. This milkweed can persist year-round in mild climates, allowing monarchs to breed throughout the winter rather than going into diapause. Tropical milkweed may foster higher loads of a monarch parasite called Oe (*Ophryocystis elektroscirrha*), which negatively impacts monarch health. Because of these implications, we recommend planting native species of milkweeds in areas where they historically occurred. You can read more about Oe in a fact sheet by the Monarch Joint Venture: http://monarchjointventure.org/images/uploads/documents/Oe_fact_sheet.pdf.

Protect Monarchs from Pesticides


Both insecticides and herbicides can be harmful to monarchs. Herbicides can reduce floral resources and host plants. Although dependent on timing, rate, and method of application, most insecticides have the potential to poison or kill monarchs and other pollinators. Systemic insecticides, including neonicotinoids, have received significant attention for their potential role in pollinator declines (imidacloprid, dinotefuran, clothianidin, and thiamethoxam are examples of systemic insecticides now found in various farm and garden products). Because plants absorb systemic insecticides as they grow, the chemicals become distributed throughout all plant tissues, including the leaves and nectar. New research has demonstrated that some neonicotinoids are toxic to monarch caterpillars that are poisoned as they feed on leaf tissue of treated plants. You can help protect monarchs by avoiding the use of these and other insecticides. Before purchasing plants from nurseries and garden centers, be sure to ask whether they have been treated with systemic insecticides. To read more about threats to pollinators from pesticides, please visit: www.xerces.org/pesticides.

Additional Resources

Gardening for Butterflies



Attracting Birds, Butterflies, and Other Backyard Wildlife



Available through www.xerces.org/books and http://bit.ly/1Xhxfgu.

Conservation Status and Ecology of the Monarch Butterfly in the U.S. Report

www.xerces.org/us-monarch-consv-report

Guide to Milkweeds and Monarchs in the Western U.S. www.xerces.org/western-us-monarch-guide

Milkweed Seed Finder www.xerces.org/milkweed-seed-finder

Websites

The Xerces Society www.xerces.org/monarchs

Monarch Joint Venture www.monarchjointventure.org/resources

Natural Resources Conservation Service www.nrcs.usda.gov/monarchs

National Wildlife Federation www.nwf.org/butterflies

Citizen Science Efforts in the Rocky Mountains Region

Southwest Monarch Study www.swmonarchs.org

Xerces Society & USFWS Milkweed and Monarch Survey www.xerces.org/milkweedsurvey

Journey North www.learner.org/jnorth/monarch

Monarch Larva Monitoring Project www.mlmp.org

Project Monarch Health www.monarchparasites.org

Acknowledgements

Nectaring data and observations, background information, and other contributions to this publication were taken from the published literature and generously provided by multiple researchers, gardeners, partners, and biologists. For the full list of data sources, please visit our website: www.xerces.org/monarch-nectar-plants. Funding provided by the Monarch Joint Venture and USDA Natural Resources Conservation Service. Additional support comes from Cascadian Farm, Ceres Trust, Cheerios, CS Fund, Disney Conservation Fund, The Dudley Foundation, The Edward Gorey Charitable Trust, General Mills, National Co+op Grocers, Nature Valley, The New-Land Foundation, Inc., Turner Foundation, Inc., Whole Foods Market and its vendors, and Xerces Society Members.

Written by Candace Fallon, Nancy Lee Adamson, Sarina Jepsen, and Mace Vaughan. Designed by Kaitlyn Rich. Formatted by Michele Blackburn. PHOTO CREDITS: Tom Potterfield*: (left cover). Peter Gorman*: 1. Justin Meissen*: 2. Andrey Zharkikh*: 3. Keir Morse***: 4. freejinn****: 5. Alejandro Bayer Tamayo*: 6. Ben VanderWeide***: 7. © Smithsonian Institution, National Museum of Natural History, Department of Botany: 8. Luigi Rignanese***: 9. Judith Lopez Sikora***: 10. Joshua Mayer*: 11, 13 (cover). © 2004 Robert Sivinski***: 12. © Scott Namestnik: 14. © 2012 Barry Rice***: 15. Brett Whaley*: 16. Kelly O'Donnell, Urban Flora of NYC: 17. Bryant Olsen*: 18. stonebird*: 19. Forest & Kim Star*: 20. Thayne Tuason*: 21. Peter Stevens*: 22 (cover). Stan Shebs**: 23. Matt Lavin*: 24. *Courtesy of flickr.com/**Wikimedia Commons/***CalPhotos/****iNaturalist. Photographs remain under the copyright of the photographer. The Xerces Society is an equal opportunity employer and provider. © 2016 by The Xerces Society for Invertebrate Conservation.

This material is based upon work supported by the Natural Resources Conservation Service, U.S. Department of Agriculture, under number 65-7482-15-118. Any opinions, findings, conclusions, or recommendations expressed in this publication are those of the author(s) and do not necessarily reflect the views of the U.S. Department of Agriculture.